

# OSWESTRY LOW BACK DISABILITY QUESTIONNAIRE

Instructions: this questionnaire has been designed to give us information as to how your back pain has affected your ability to manage everyday life. Please answer every section and mark in each section only the ONE box which applies to you at this time. We realize you may consider 2 of the statements in any section may relate to you, but please mark the box which most closely describes your current condition.

## 1. PAIN INTENSITY

- ☐ I can tolerate the pain I have without having to use pain killers
- ☐ The pain is bad but I manage without taking pain killers
- ☐ Pain killers give complete relief from pain
- ☐ Pain killers give moderate relief from pain
- ☐ Pain killers give very little relief from pain
- ☐ Pain killers have no effect on the pain and I do not use them

## 2. PERSONAL CARE (e.g. Washing, Dressing)

- ☐ I can look after myself normally without causing extra pain
- ☐ I can look after myself normally but it causes extra pain
- ☐ It is painful to look after myself and I am slow and careful
- ☐ I need some help but manage most of my personal care
- ☐ I need help every day in most aspects of self care
- ☐ I don't get dressed, I was with difficulty and stay in bed

## 3. LIFTING

- ☐ I can lift heavy weights without extra pain
- ☐ I can lift heavy weights but it gives extra pain
- ☐ Pain prevents me from lifting heavy weights off the floor, but I can manage if they are conveniently positioned, i.e. on a table
- ☐ Pain prevents me from lifting heavy weights, but I can manage light to medium weights if they are conveniently positioned
- ☐ I can lift very light weights
- ☐ I cannot lift or carry anything at all

## 4. WALKING

- ☐ Pain does not prevent me walking any distance
- ☐ Pain prevents me walking more than one mile
- ☐ Pain prevents me walking more than ½ mile
- ☐ Pain prevents me walking more than ¼ mile
- ☐ I can only walk using a stick or crutches
- ☐ I am in bed most of the time and have to crawl to the toilet

## 5. SITTING

- ☐ I can sit in any chair as long as I like
- ☐ I can only sit in my favorite chair as long as I like
- ☐ Pain prevents me from sitting more than one hour
- ☐ Pain prevents me from sitting more than ½ hour
- ☐ Pain prevents me from sitting more than 10 minutes
- ☐ Pain prevents me from sitting at all

## 6. STANDING

- ☐ I can stand as long as I want without extra pain
- ☐ I can stand as long as I want but it gives me extra pain
- ☐ Pain prevents me from standing for more than one hour
- ☐ Pain prevents me from standing for more than 30 minutes
- ☐ Pain prevents me from standing for more than 10 minutes
- ☐ Pain prevents me from standing at all

## 7. SLEEPING

- ☐ Pain does not prevent me from sleeping well
- ☐ I can sleep well only by using medication
- ☐ Even when I take medication, I have less than 6 hrs sleep
- ☐ Even when I take medication, I have less than 4 hrs sleep
- ☐ Even when I take medication, I have less than 2 hrs sleep
- ☐ Pain prevents me from sleeping at all

## 8. SOCIAL LIFE

- ☐ My social life is normal and gives me no extra pain
- ☐ My social life is normal but increases the degree of pain
- ☐ Pain has no significant effect on my social life apart from limiting my more energetic interests, i.e. dancing, etc.
- ☐ Pain has restricted my social life and I do not go out as often
- ☐ Pain has restricted my social life to my home
- ☐ I have no social life because of pain

## 9. TRAVELLING

- ☐ I can travel anywhere without extra pain
- ☐ I can travel anywhere but it gives me extra pain
- ☐ Pain is bad, but I manage journeys over 2 hours
- ☐ Pain restricts me to journeys of less than 1 hour
- ☐ Pain restricts me to short necessary journeys under 30 minutes
- ☐ Pain prevents me from traveling except to the doctor or hospital

## 10. EMPLOYMENT/ HOME MAKING

- ☐ My normal homemaking/ job activities do not cause pain.
- ☐ My normal homemaking/ job activities increase my pain, but I can still perform all that is required of me.
- ☐ I can perform most of my homemaking/ job duties, but pain prevents me from performing more physically stressful activities (e.g. lifting, vacuuming)
- ☐ Pain prevents me from doing anything but light duties.
- ☐ Pain prevents me from doing even light duties.
- ☐ Pain prevents me from performing any job or homemaking chores.

Patient name: \_\_\_\_\_

Date: \_\_\_\_\_

# Opioid Safety Survey

Because there is addiction risk with opioid medicines, we must first understand your history before we set a treatment plan for you.

Please circle "Yes" or "No" in the chart below as the case applies to you today or in the past. Then give this sheet to your doctor or nurse.

Patient Name: \_\_\_\_\_

Date: \_\_\_\_\_

Mark each box that applies	Female	Male
<b>Do YOU have a history of substance abuse of any of the following?</b>		
Alcohol	YES / NO	YES / NO
Illegal drugs	YES / NO	YES / NO
Prescription drugs	YES / NO	YES / NO
<b>Do you have a FAMILY history of substance abuse of any of the following?</b>		
Alcohol	YES / NO	YES / NO
Illegal drugs	YES / NO	YES / NO
Prescription drugs	YES / NO	YES / NO
<b>Are you between 16—45 years old?</b>		
	YES / NO	YES / NO
<b>Were you sexually abused as a child?</b>		
	YES / NO	YES / NO
<b>Have you had one of the following mental health conditions?</b>		
ADD, OCD, bipolar, schizophrenia	YES / NO	YES / NO
Depression	YES / NO	YES / NO

Patient Name: \_\_\_\_\_ Date: \_\_\_\_\_

## PATIENT HEALTH QUESTIONNAIRE-9 (PHQ-9)

Over the last 2 weeks, how often have you been bothered  
by any of the following problems?  
(Use "✓" to indicate your answer)

	Not at all	Several days	More than half the days	Nearly every day
1. Little interest or pleasure in doing things	0	1	2	3
2. Feeling down, depressed, or hopeless	0	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having little energy	0	1	2	3
5. Poor appetite or overeating	0	1	2	3
6. Feeling bad about yourself — or that you are a failure or have let yourself or your family down	0	1	2	3
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
8. Moving or speaking so slowly that other people could have noticed? Or the opposite — being so fidgety or restless that you have been moving around a lot more than usual	0	1	2	3
9. Thoughts that you would be better off dead or of hurting yourself in some way	0	1	2	3

FOR OFFICE CODING 0 + \_\_\_\_\_ + \_\_\_\_\_ + \_\_\_\_\_  
=Total Score: \_\_\_\_\_

If you checked off any problems, how difficult have these problems made it for you to do your  
work, take care of things at home, or get along with other people?

Not difficult at all <input type="checkbox"/>	Somewhat difficult <input type="checkbox"/>	Very difficult <input type="checkbox"/>	Extremely difficult <input type="checkbox"/>
---	---	---	--

# FRAT

Date .....

Name .....

Score 1 for every category and total at the bottom of the 2 columns		Yes	No
1	Is there a history of any fall in the previous year?		
2	Is the client on four or more medications per day?		
3	Does the client have a diagnosis of stroke or Parkinson's?		
4	Does the client report any problems with their balance?		
5	Is the client <b>unable</b> to rise from a chair of knee height without using their arms?		
<b>Total</b>			

## Level of predicted risk:

### 3 - 5 yes's = higher falls risk

Complete full falls risk assessment (see FESI tool or service user risk assessment)

Consult health and social care professionals.

Complete provider self assessment.

### Less than 3 yes's = lower risk

If the person has mobility problems consider referral to Community Therapy Services

Carry out exercises with the individual