

# **Colonoscopy with Oral Lavage Prep**

Patient Name:	
Date of Procedure	
Arrival Time	
Procedure Time	

820 Bestgate Road Suite 1A Annapolis, Maryland 21401 410-224-2116

## Your Procedure:

You are scheduled for a colonoscopy. The doctor will examine the lining of your large intestine with a colonoscope. If polyps or other abnormalities are found during the procedure, the doctor will remove the abnormal tissue for further evaluation.

Maryland Center for Digestive Health
820 Bestgate Road, Suite 1A
Annapolis, Maryland 21401

**Baltimore Washington Medical Center** 

Place:

1 Week Prior	5 Days Prior	2 Days Prior	1 Day Prior	Procedure Day
Purchase the following: - Bisacodyl tablets Colyte, Golyte,Trilyte, Nulytely from your pharmacy.	Begin low fiber diet, see reverse.	Continue low fiber diet.  Drink at least four 8oz glasses of water throughout	Begin clear liquid diet. No solid foods today. Red, purple or orange colored liquids are not allowed. You may have small	6 hours prior to procedure: Drink the remaining half of the prep solution. Drink 8 ounces every 15
Follow these instructions, not the box instructions.	Stop consuming all high-fiber foods.	the day.	amounts of coffee or tea without cream.	minutes until finished. This should take 2 hours.
If you take dietary, herbal or fiber supplements or medications			Drink at least four 8oz glasses of water throughout the day.	You may take your essential
containing iron, including multi- vitamins with iron, discontinue these 7 days before your				morning medications including Blood Pressure Medication with a few small sips of water,
appointment.	If you are taking ibuprofen (Advil, Nuprin or Motrin) or other non-steroidal (NSAID)	No solid foods	2:00 pm Take Bisacodyl 10 mg.	at least 4 hours prior to your procedure, unless otherwise directed by your physician.
lf you are taking blood thinners including Coumadin (Warfarin), Pradaxa (Dibigatron), Xarelto	anti-inflammatory medications, <u>stop taking</u> them today. Tylenol and	after midnight!	5:00 pm Thoroughly mix 4 liters (1 gallon) clear liquid to the Colyte, Golyte,Trilyte, Nulytely powder.	NOTHING by mouth of
(Rivaroxaban), Effient (Prasugrel) Plavix (Clopidogrel) Brilinta (Ticagrelor) and Eliquis	Celebrex are okay to use.		Do Not Refrigerate.	NOTHING by mouth at least 4 hours prior to your scheduled procedure time
(Apixaban), you will need an office visit prior to your procedure.			<b>6pm:</b> Drink 8 ounces of the prep solution and continue drinking an	- including gum, tobacco, hard candy/mints!
You may <u>continue</u> your "baby" aspirin and regular aspirin if you were			8 ounce glass of solution every 15 minutes <i>until half of the solution is finished.</i> This should take	You will need:
prescribed them for a cardiovascular condition.			about 2 hours. Remaining solution to be taken 6 hours prior to your procedure time.	-Insurance Cards -Driver's License/Photo ID
Hold Subcutaneous Injection 1 week prior to procedure.			If you experience nausea or	<ul> <li>-Any co-insurance fees due</li> <li>-A responsible adult driver to drive you home</li> </ul>
Ozempic, Wegovy, Trulicity, Bydureon, Mounjaro - ususally weekly injections			vomiting take a 15-30 minute break, then continue drinking prep solution.	- Referral

Phone: 410-224-2116 Fax: 410.224.2118

## Milk and dairy





Meat



### Nuts, nut butter, seeds



Fats and oils



Soups



### OK to eat:

- · Milk
- · Cream
- · Hot chocolate
- Buttermilk
- · Cheese, including cottage cheese
- · Yogurt
- · Sour cream

## NO yogurt mixed with:

- · nuts, seeds, granola
- · fruit with skin or seeds (such as berries)

## Legumes





## None allowed

### NO:

- · Dried peas (including split or black-
- · Dried beans (including kidney, pinto, garbanzo/chickpea)
  - Lentils
  - · Any other legume

## OK to eat:

- · Breads and grains made with refined white flour (including rolls, muffins, bagels, pasta)
- · White rice

OK to eat:

Chicken

Turkey

• Lamb

Veal

Eggs

Tofu

· Lean pork

· Fish and seafood

- · Plain crackers, such as Saltines
- · Low-fiber cereal (including puffed rice, cream of wheat, corn flakes)

## NO whole grains or high-fiber:

- . Brown or wild rice
- · Whole grain bread, rolls, pasta, or crackers
- · Whole grain or high-fiber cereal (including granola, raisin bran, oatmeal)
- . Bread or cereal with nuts or seeds

## Fruits







## OK to eat:

- . Fruit juice without pulp
- Applesauce
- · Ripe cantaloupe and honeydew
- · Ripe, peeled apricots and peaches
- · Canned or cooked fruit without seeds or skin

### NO seeds, skin, membranes; or dried fruit:

- . Raw fruit with seeds, skin, or membranes (includes berries, pine apple, apples, oranges, watermelon)
- · Any cooked or canned fruit with seeds or skin
- · Raisins or other dried fruit

## NO tough meat with gristle

## Vegetables



## OK for some if cooked or canned:

- · Canned or cooked vegetables without skin or peel (includes peeled carrots, mushrooms, turnips, asparagus tips)
- · Potatoes without skin
- · Cucumbers without seeds or peel

## NO raw, skin, seeds, peel; or certain other vegetables:

- Corn
- · Potatoes with skin
- Tomotoes
- · Cucumbers with seeds and peel
- Cooked cabbage or Brussels sprouts
- · Green peas
- . Summer and winter squash
- Lima beans
- Onions



## OK to eat:

OK to eat:

Margarine

Mayonnaise

or nuts

· Vegetable and other oils

Butter

· Creamy (smooth) peanut or almond butter

### NO nuts or seeds:

- Nuts including peanuts, almonds, walnuts
- · Chunky nut butter

No:

• Chili

Lentil soup

· Corn soup

Pea soup

Unstrained soups

· Dried bean soup

· Seeds such as fennel, sesame. pumpkin, sunflower

### Desserts





- OK to eat: Custard
- · Plain pudding
- · Ice cream
- · Sherbet or sorbet
- . Jell-O or gelatin without added fruit or red or purple dye
- . Cookies or cake made with white flour, prepared without seeds, dried fruit, or nuts

### NO:

- Coconut.
- · Anything with seeds or nuts
- · Anything with added red or purple
- · Cookies or cakes made with whole grain flour, seeds, dried fruit, or nuts

## OK to eat:

- . Broth, bouillon, consomme, and strained soups
- · Milk or cream-based soup, strained

· Salad dressings made without seeds

## or nuts



Other



### OK to eat:

- Coffee
- Ton
- · Hot chocolate or cocoa
- · Clear fruit drinks (no pulp)
- · Soda and other carbonated beverages
- · Ensure, Boost, or Enlive without added fiber

- · Fruit or vegetable juice with pulp
- · Beverages with red or purple dye

### OK to eat:

- Sugar
- · Salt Jelly
- Honey Syrup
- Lemon juice

## NO:

- Coconut
- · Popcorn
- · Jam Marmalade
- · Relishes
- Pickles Olives
- · Stone-ground mustard