

Colonoscopy with SuTab Prep

Patient Name:	
Date of Procedure	
Arrival Time	
Procedure Time	

Annapolis, Maryland 21401 410-224-2116

Your Procedure:

You are scheduled for a colonoscopy. The doctor will examine the lining of your large intestine with a colonoscope. If polyps or other abnormalities are found during the procedure, the doctor will remove the abnormal tissue for further evaluation.

820 Bestgate Road, Suite 1A Annapolis, Maryland 21401

Baltimore Washington Medical Center

Maryland Center for Digestive Health

YOUR BOWEL PREP IS EXTREMELY IMPORTANT! PLEASE READ INSTRUCTIONS CAREFULLY

Place:

1 Week Prior	5 Days Prior	2 Days Prior	1 Day Prior	Procedure Day
Have your prescription filled today.	Begin low fiber diet. See Reverse	Continue low fiber diet.	Begin clear liquid diet. No solid foods today. Red, purple or	Continue to consume only clear liquids until after the colonoscopy.
Follow these instructions, not the box instructions.	Stop consuming all high- fiber foods. This includes food with seeds and nuts.	Drink at least four 8oz glasses of water throughout the day.	orange colored liquids are not allowed. You may have small amounts of coffee or tea without cream.	• The morning of colonoscopy 6 hours prior to the colonoscopy open the second bottle of 12 tablets.
If you take dietary, herbal or fiber supplements or medications containing iron, including multivitamins with iron, discontinue	If you are taking ibuprofen		Drink at least four 8oz glasses of water throughout the day.	• Repeat STEP 1 to STEP 3 from Dose 1.
these 7 days before your If you are taking blood thinners including Coumadin (Warfarin), Pradaxa (Dibigatron), Xarelto (Rivaroxaban), Effient (Prasugrel) Plavix (Clopidogrel) Brilinta	(Advil, Nuprin or Motrin) or other non-steroidal (NSAID) anti-inflammatory medications, stop taking them today. Tylenol and Celebrex are okay to use.	No solid foods after midnight!	Step 1: 6pm: Open 1 bottle of 12 tablets and fill the provided container with 16 ounces of water (up to the fill line). Swallow each tablet every 1to 2 minutes with a sip of water, and drink the entire amount of water within 20 minutes.	You may take your essential morning medications including <u>Blood Pressure Medication</u> with a few small sips of water, at least 4 hours prior to your procedure, unless otherwise directed by your physician.
(Ticagrelor) and Eliquis (Apixaban), you will need an office visit prior to your procedure. You may continue your "baby" aspirin and regular aspirin if you were prescribed them for a cardiovascular condition.			Step 2:Approximately 1 hour after the last tablet is ingested, fill the provided container again with 16 ounces of water (up to the fill line), and drink the entire amount over 30 minutes. Step 3: Approximately 30 minutes after finishing the second container of water,	
Hold Subcutaneous Injection 1 week prior to procedure. Ozempic, Wegovy, Trulicity, Bydureon, Mounjaro - ususally weekly injections			fill the provided container with 16 ounces of water (up to the fill line), and drink the entire amount over 30 minutes. If you experience nausea or vomiting take a 15-30 minute break, then continue drinking prep solution.	You will need: -Insurance Cards -Driver's License/Photo ID -Any co-insurance fees due -A responsible adult driver to drive you home -Referral

You will receive your procedure results via the portal. Visit <u>www.aagastro.com</u> to sign up today.

Phone: 410-224-2116 Fax: 410.224.2118

Milk and dairy







Nuts, nut butter, seeds



Fats and oils



Soups



OK to eat:

- · Milk
- · Cream
- · Hot chocolate
- Buttermilk
- · Cheese, including cottage cheese

. Breads and grains made with refined

white flour (including rolls, muffins,

· Plain crackers, such as Saltines

· Low-fiber cereal (including puffed

rice, cream of wheat, corn flakes)

· Yogurt

OK to eat:

bagels, pasta)

· White rice

· Sour cream

NO yogurt mixed with:

- · nuts, seeds, granola
- · fruit with skin or seeds (such as berries)

NO whole grains or high-fiber:

. Whole grain bread, rolls, pasta,

· Whole grain or high-fiber cereal

(including granola, raisin bran,

· Bread or cereal with nuts or seeds

NO tough meat with gristle

. Brown or wild rice

or crackers

oatmeal)

Legumes





Fruits



. Fruit juice without pulp Applesauce

seeds or skin

OK to eat:

None allowed

- · Ripe cantaloupe and honeydew
- · Ripe, peeled apricots and peaches · Canned or cooked fruit without

OK for some if cooked or canned:

without skin or peel (includes peeled

· Canned or cooked vegetables

carrots, mushrooms, turnips,

· Cucumbers without seeds or peel

. Jell-O or gelatin without added fruit

. Cookies or cake made with white flour, prepared without seeds, dried

asparagus tips)

· Potatoes without skin

NO seeds, skin, membranes; or dried fruit:

garbanzo/chickpea)

· Any other legume

· Raw fruit with seeds, skin, or membranes (includes berries, pine apple, apples, oranges, watermelon)

· Dried peas (including split or black-

· Dried beans (including kidney, pinto,

eyed)

• Lentils

- · Any cooked or canned fruit with seeds or skin
- . Raisins or other dried fruit

Meat



Eggs Tofu

OK to eat:

Margarine

Mayonnaise

or nuts

OK to eat:

strained soups

· Vegetable and other oils

· Salad dressings made without seeds

. Broth, bouillon, consomme, and

· Milk or cream-based soup, strained

Butter

OK to eat:

Chicken

Turkey

• Lamb

Veal

· Lean pork

OK to eat:

Fish and seafood

· Creamy (smooth) peanut or almond butter

NO nuts or seeds:

- · Nuts including peanuts, almonds, walnuts
- · Chunky nut butter

or nuts

Unstrained soups

· Dried bean soup

• Chili

Lentil soup

· Corn soup

· Pea soup

· Seeds such as fennel, sesame. pumpkin, sunflower

Vegetables







Desserts





Other

NO salad dressing made with seeds Drinks or beverages



OK to eat:

- Coffee

- · Hot chocolate or cocoa
- . Clear fruit drinks (no pulp)
- beverages
- . Ensure, Boost, or Enlive without added fiber

NO raw, skin, seeds, peel; or certain

- other vegetables: · Com
- · Potatoes with skin
- Tomatoes
- · Cucumbers with seeds and peel
- Cooked cabbage or Brussels sprouts
- · Green peas
- . Summer and winter squash
- Lima beans
- Onions

NO:

- · Coconut
- · Anything with seeds or nuts
- · Anything with added red or purple
- · Cookies or cakes made with whole grain flour, seeds, dried fruit, or nuts

- Tea

OK to eat:

Custard

· Ice cream

· Plain pudding

fruit, or nuts

· Sherbet or sorbet

or red or purple dye

- · Soda and other carbonated

- · Fruit or vegetable juice with pulp
- . Beverages with red or purple dye

OK to eat: • Sugar

- Salt
- · Jelly Honey
- Syrup
- · Lemon juice

NO:

- Coconut
- · Popcorn
- Jam
- Marmalade Relishes
- Pickles
- Olives
- · Stone-ground mustard