



**ANNE ARUNDEL
GASTROENTEROLOGY
ASSOCIATES, P.A.**

Christopher F. Schultz, MD
Melanie L. Jackson, MD, FACP
Praveena G. Velamati, MD
Christopher A. Olenec, MD
Thomas L. Simcox, MD
Raja Taunk, MD
Mark Flasar, MD, MS
David S. Jencks, MD
Raza Hasan, MD
Suresh Misra, MD
Amanda M. Era, CRNP
Lauren Brennan, CRNP
Emily Nehme, PA
Makeda Berhane, CRNP

Bestgate Medical Clinic 820
Bestgate Road, Suite 2A
Annapolis, Maryland 21401

Kent Island
200 Saint Claire Place Suite
100, Stevensville, MD 21666

Odenton
1215 Annapolis Road Suite
201, Odenton, MD 21113

Bowie
4175 North Hanson Court Suite
304, Bowie, MD 20716

Pasadena
8109 Ritchie Highway Suite
102, Pasadena, MD 21122

Greenbelt
7500 Greenway Center Drive
Suite 620, Greenbelt, MD
20770



410-224-2116

Fax 410-224-2118

www.aagastro.com

Toll Free 1-866-224-AAGA

Local 410-280-AAGA (2242)

DIABETIC INSTRUCTIONS FOR COLONOSCOPY FOR DIABETIC PATIENTS INSULIN DEPENDENT / ORAL HYPOGLYCEMICS

Hold Subcutaneous Injection 1 week prior to procedure. Ozempic, Wegovy, Trulicity, Bydureon, Mounjaro - usually weekly injections

MORNING COLONOSCOPY

If your Insulin is managed on a sliding scale by your Endocrinologist or PCP, follow their recommendations. If questions, consult your Endocrinologist or PCP.

If you do not use a sliding scale regimen, use the following procedure:

INSULIN DEPENDENT

- Take 1/2 usual dose day prior to procedure. (if you take insulin twice a day, take 1/2 usual dose both times)
- Do prep
- Do finger sticks
- Take 1/2 usual dose before your procedure
- When you have eaten after the procedure, take 1/2 your usual dose of insulin

ORAL DIABETIC MEDICATION

- None the day prior to procedure
- Do prep
- None the day of the procedure
- When you have eaten after the procedure, resume your usual dose

PATIENT RESPONSIBILITY

IT IS YOUR RESPONSIBILITY TO CHECK WITH YOUR INSURANCE COMPANY REGARDING PRE-AUTHORIZATION AND ANY REFERRALS NECESSARY FOR THIS PROCEDURE. YOU WILL NEED TO PRESENT A CURRENT PHOTO ID, VALID INSURANCE CARD AND A VALID REFERRAL, IF REQUIRED. YOU WILL BE EXPECTED TO PAY ANY COPAYMENTS, COINSURANCE, DEDUCTIBLES AND OUTSTANDING BALANCES AT THE TIME OF YOUR APPOINTMENT. IF YOU ARE NOT PREPARED TO PAY THE FEES DUE, YOUR APPOINTMENT MAY BE CANCELLED.

PLEASE LEAVE VALUABLES AT HOME OR WITH THE RESPONSIBLE TRANSPORTATION PROVIDER. PLEASE BRING YOUR EYEGLASSES, IF APPLICABLE, AS WELL AS SUFFICIENT CASH, CREDIT CARD OR CHECK FOR YOUR FINANCIAL RESPONSIBILITY.

IT WILL BE NECESSARY FOR YOUR TRANSPORTATION PROVIDER TO WAIT FOR YOU.

Anne Arundel Gastroenterology Associates provides medical services regardless of race, color, age, national origin, sex, religion, or handicap. If you are hearing impaired or have a language barrier, please notify our office.

DIABETIC INSTRUCTIONS FOR COLONOSCOPY FOR DIABETIC PATIENTS INSULIN DEPENDENT/ORAL HYPOGLYCEMICS

Hold Subcutaneous Injection 1 week prior to procedure. Ozempic, Wegovy, Trulicity, Bydureon, Mounjaro - usually weekly injections

AFTERNOON COLONOSCOPY

If your insulin is managed on a sliding scale by your Endocrinologist or PCP, follow their recommendations. If questions, consult your Endocrinologist or PCP.

If you do not use a sliding scale regimen, use the following procedure:

INSULIN DEPENDENT

- Take 1/2 usual dose day prior to procedure. (if you take insulin twice a day, take 1/2 usual dose both times)
- Take 1/2 usual dose before your procedure
- Do finger sticks
- When you have eaten after the procedure, take 1/2 your usual dose if you normally take insulin twice a day

ORAL DIABETIC MEDICATION

- None day prior to procedure
- None day of procedure
- Do prep
- When you have eaten after the procedure, and you normally do not take the p.m. dose, take 1/2 your usual dose; otherwise resume your usual dose.