

Colonoscopy with 2day Miralax

| Date of Procedure | |
|-------------------|--|
| Arrival Time | |

Patient Name:

Procedure Time

Place:

820 Bestgate Road Suite 1A Annapolis, Maryland 21401 410-224-2116

Your Procedure:

You are scheduled for a colonoscopy. The doctor will examine the lining of your large intestine with a colonoscope. If polyps or other abnormalities are found during the procedure, the doctor will remove the abnormal tissue for further evaluation.

| Maryland Center for Digestive Health |
|--|
| 820 Bestgate Road, Suite 1A Annapolis, |
| Maryland 21401 |

□ Baltimore Washington Medical Center

| 1 Week Prior | 5 Days Prior | 2 Days Prior | 1 Day Prior | Procedure Day |
|---|---|---|---|---|
| Purchase the Follow: Items listed below are over the counter NOT a Prescription. Follow these instructions, not the box instructions. | Begin low fiber diet. Stop consuming all high-fiber foods. This includes food with seeds and nuts. | Clear Liquids only. Drink at least four 8oz glasses of water throughout the day. | Begin clear liquid diet. No solid foods today. Red, purple or orange colored liquids are not allowed. You may have small amounts of coffee or tea without cream | 6 hours prior to procedure: Take the second 1/2 of the Gatorade/Miralax solution Drink one 8oz glass every 15 minutes until completed |
| -Dulcolax tablets, six 5mg (not suppositories) -(2 bottles) 64oz Gatorade or PowerAde: (no red orange or blue) -(2 bottles)Miralax 238gm If you take dietary, herbal or fiber supplements or medications containing iron, including multivitamins with iron, discontinue these 7 days before your appointment. If you are taking blood thinners including Coumadin (Warfarin), Pradaxa (Dibigatron), Xarelto (Rivaroxaban), Effient (Prasugrel) Plavix (Clopidogrel) Brilinta (Ticagrelor) and Eliquis (Apixaban), you will need an office visit prior to your procedure. You may continue your "baby" aspirin and regular aspirin if you were prescribed them for a Hold Subcutaneous Injection 1 week prior to procedure. Ozempic, Wegovy, Trulicity, Bydureon, Mounjaro ususally weekly injections | If you are taking ibuprofen (Advil, Nuprin or Motrin) or other non-steroidal (NSAID) anti-inflammatory medications, stop taking them today. Tylenol and Celebrex are okay to use. 3 days prior No solid foods after midnight! | 5:00pm Thoroughly mix the entire bottle of Miralax in the bottle of clear Gatorade or PowerAde, (you may want to pour 8oz out of the bottle before mixing) Do not refrigerate longer than 1 hour before starting your prep. Do not over chill. 6:00pm Drink ½ of the solution 8:00pm Drink ½ of the miralax/Gatorade solution then Drink one 8oz glass and repeat every 15miutes until you finish the 32ounces. | Drink at least four 8oz glasses of Water throughout the day. | You may take your essential morning medications including Blood Pressure Medication with a few small sips of water, at least 4 hours prior to your procedure, unless otherwise directed by your physician. NOTHING by mouth at least 4 hours prior to your scheduled procedure time – including gum, tobacco, hard candy/mints! You will need: -Insurance Cards -Driver's License/Photo ID -Any co-insurance fees due -A responsible adult driver to drive you home -Referral |